

IN THE SPECIFICATION

Please make the amendments to the Specification as indicated below:

Please replace the two paragraphs on page 5, lines 3-22 with the following two paragraphs:

MN/CA IX has a number of properties that distinguish it from other known CA isoenzymes and evince its relevance to oncogenesis. Those properties include its density dependent expression in cell culture (e.g., HeLa cells), its correlation with the tumorigenic phenotype of somatic cell hybrids between HeLa and normal human fibroblasts, its close association with several human carcinomas and its absence from corresponding normal tissues [e.g., Zavada et al., Int. J. Cancer, 54: 268-274 (1993); Pastorekova et al., Virology, 187: 620-626 (1992); Liao et al., Am. J. Pathol., 145: 598-609 (1994); Pastorek et al., Oncogene, 9: 2788-2888 (1994); Cote, Women's Health Weekly: News Section, p. 7 (March 30, 1998); Liao et al., Cancer Res., 57: 2827 (1997); Vermylen et al., "Expression of the MN antigen as a biomarker of lung carcinoma and associated precancerous conditions," Proceedings AACR, 39: 334 (1998); McKiernan et al., Cancer Res., 57: 2362 (1997); and Turner et al., Hum. Pathol., 28(6): 740 (1997)]. In addition, the *in vitro* transformation

potential of MN/CA IX cDNA has been demonstrated in NIH 3T3 fibroblasts [Pastorek et al., id.].

AP ✓
The MN protein has also been identified with the G250 antigen. Uemura et al., "Expression of Tumor-Associated Antigen MN/G250 in Urologic Carcinoma: Potential Therapeutic Target, " J. Urol., 154 (4 Suppl.): 377 (Abstract 1475; 1997) states:
"Sequence analysis and database searching revealed that G250 antigen is identical to MN, a human tumor-associated antigen identified in cervical carcinoma (Pastorek et al., 1994)."

Please replace Table 1 on page 26 with the following new Table 1:

TABLE 1

Exon-Intron Structure of the Human MN Gene

Exon	Size	Genomic Position**	SEQ ID NO	5'splice acceptor	SEQ ID NO
1	445	*3507-3951	28	AGAAG gtaagt	67
2	30	5126-5155	29	TGGAG gtgaga	68
3	171	5349-5519	30	CAGTC gtgagg	69
4	143	5651-5793	31	CCGAG gtgagc	70
5	93	5883-5975	32	TGGAG gtacca	71
6	67	7376-7442	33	GGAAG gtcagt	72
7	158	8777-8934	34	AGCAG gtgggc	73
8	145	9447-9591	35	GCCAG gtacag	74
9	27	9706-9732	36	TGCTG gtgagt	75
10	82	10350-70431	37	CACAG gtatta	76
11	191	10562-10752	38	ATAAT end	
Intron	Size	Genomic Position **	SEQ ID NO	3'splice acceptor	SEQ ID NO
1	1174	3952-5125	39	atacag GGGAT	77
2	193	5156-5348	40	ccccag GCGAC	78
3	131	5520-5650	41	acgcag TGCAA	79
4	89	5794-5882	42	tttcag ATCCA	80
5	1400	5976-7375	43	ccccag GAGGG	81
6	1334	7443-8776	44	tcacag GCTCA	82
7	512	8935-9446	45	ccctag CTCCA	83
8	114	9592-9705	46	ctccag TCCAG	84
9	617	9733-10349	47	tcgcag GTGACA	85
10	130	10432-10561	48	acacag AAGGG	86

** positions are related to nt numbering in whole genomic sequence including the 5' flanking region [Figure 2A-F]

* number corresponds to transcription initiation site determined below by RNase protection assay